

LASCAR GROUPS AND THE FIRST HOMOLOGY GROUPS OF STRONG TYPES IN ROSY THEORIES

JUNGUK LEE

ABSTRACT. For a rosy theory, we give a canonical surjective homomorphism from a Lascar group over $A = \text{acl}^{eq}(A)$ to a first homology group of a strong type over A , and we describe its kernel by an invariant equivalence relation. As a consequence, we show that the first homology groups of strong types in rosy theories have the cardinalities of one or at least $2^{\aleph_0}$. We give two examples of rosy theories having non trivial first homology groups of strong types over $\text{acl}^{eq}(\emptyset)$. In these examples, these two homology groups are exactly isomorphic to their Lascar group over $\text{acl}^{eq}(\emptyset)$.

1. INTRODUCTION

In model theory, strong types and Lascar types are important objects to understand invariant equivalence relations in a theory. Let $T = T^{eq}$ be a theory and $\mathcal{C}$ be a monster model of T . Let A be a small set in $\mathcal{C}$. We say that an equivalence relation E is *finite* if it has finitely many E -classes, *bounded* if the number of E -classes are less than the cardinality of $\mathcal{C}$, and *A -invariant* if for a, b and $f \in \text{Aut}_A(\mathcal{C})$, $(a, b) \in E$ implies $(f(a), f(b)) \in E$. We say finite tuples a, b have the same *strong(or Shelah) type over A* , written $a \equiv_A^s b$ if for any A -definable finite equivalence relation E over $\mathcal{C}^{[a]}$, $(a, b) \in E$. It is well known that $a \equiv_A^s b$ if and only if $a \equiv_{\text{acl}(A)} b$. Thus we take $\text{stp}(a/A)$, the strong type of a over A as $\text{tp}(a/\text{acl}(A))$, and it is the orbit of a under the action of $\text{Aut}_{\text{acl}(A)}(\mathcal{C})$. So, it makes sense to consider a strong type of an infinite tuple over A . The Lascar types over A are defined as follows : We say tuples a, b possibly infinite length have the same *Lascar type over A* , written $a \equiv_A^L b$ if for any A -invariant bounded equivalence relation E over $\mathcal{C}^{[a]}$, $(a, b) \in E$. Like strong types, a well known fact is that $a \equiv_A^L b$ if and only if there is a finite sequence $(a_0, a_1, \dots, a_n)$ in $\mathcal{C}^{[a]}$ such that $a_0 = a$, $a_n = b$, and $a_i \equiv_{M_i} a_{i+1}$ for a submodel M_i of $\mathcal{C}$ containing A for each $i = 0, \dots, n-1$. So $\text{Ltp}(a/A)$, the Lascar type of a over A is the orbit of a under the action of $\text{Autf}_A(\mathcal{C})$ which is the group of automorphism fixing a submodel containing A , and $\text{Autf}_A(\mathcal{C})$ is a normal subgroup in $\text{Aut}_A(\mathcal{C})$. So, it is well defined the

group $\text{Gal}_L(\mathcal{C}; A) = \text{Aut}_A(\mathcal{C}) / \text{Aut}_f_A(\mathcal{C})$, called the Lascar group over A . One asks when $\text{stp} = \text{Ltp}$ in T and it is determined by $\text{Gal}_L(\mathcal{C}; A)$ for $A = \text{acl}(A)$.

Two notions of strong and Lascar types are also important in the context of classification theory. In stable theories, each type over a model A is stationary; it has only one of unique non-forking extension over each B containing A . And this stationarity over a model is a characterization property of stable theories. In the case when a theory has a ternary invariant independence relation satisfying symmetry, transitivity, extension, finite and local characters, and stationarity over a model, it is well known that the theory is stable. In simple theories, the stationarity over models is substituted by type amalgamation over a model (or 3-amalgamation); for a model A and a_1, a_2, b_1, b_2 with $a_1 \equiv_A a_2$, if $\{a_i, b_i\}$ for $i = 1, 2$ and $\{b_1, b_2\}$ are independent over A , then there is $a_3 \equiv_{Ab_i} a_i$ for $i = 1, 2$ such that $\{a_3, b_1, b_2\}$ is independent over A . One generalizes 3-amalgamation to n -amalgamation for $n \geq 3$, called generalized amalgamation properties. We assume $T = T^{\text{heq}}$ in the case that T is simple. It is well known that each strong type over A which is a type over $\text{acl}(A)$ is stationary in a stable theory T , and if T is simple, $a \equiv_A^L b$ if and only if $a \equiv_{\text{bdd}(A)} b$ and each types over $\text{bdd}(A)$ satisfies 3-amalgamation, where $\text{bdd}(A)$ is the set of elements in $\mathcal{C}^1$ whose the cardinality of orbit under $\text{Aut}_A(\mathcal{C})$ is less than $|\mathcal{C}|$. In [4], J. Goodrick, B. Kim, and A. Kolesnikov introduced homology groups related with generalized amalgamation property for strong types in the context of rosy theories. They computed the first homology groups for some cases in [4] (which were all zero), and in stable theories (of course, 3-amalgamation holds in this case), they gave an explicit description of higher homology groups in [3][5]. It was not known much about the first homology groups in general. We supposed that the first homology group for a strong type p in a rosy theory is related with a Lascar group. Indeed, in [8], B. Kim, S. Kim, and the author showed that for a Lascar strong type p , the first homology group for p is always zero, so even though in a rosy theory, Lascar strong type has no 3-amalgamation still it satisfies a kind of complicated form of amalgamation.

In this paper, instead of the original definition of the first homology group in [4], we work with a restricted one, called a first homology group *over* A in a strong type over A for $A = \text{acl}(A)$ and we give a canonical surjective homomorphism from the Lascar group over A into these first homology groups in strong types over A in rosy theories. We describe its kernel by an invariant bounded equivalence relation whose classes are described by some subgroup of the automorphism group.

From this description of the kernel, we deduce that the cardinality of this first homology group for a strong type is always one or at least $2^{\aleph_0}$. At last, we give two examples of rosy theories having a strong type over $\text{acl}^{eq}(\emptyset)$ with a non trivial first homology group exactly isomorphic to their Lascar groups over $\text{acl}^{eq}(\emptyset)$. From known examples in [4][8] and our two examples, we conjecture that these first homology group in strong types over A are isomorphic to the abelianization of Lascar group over $A = \text{acl}(A)$ under the assumption that the algebraic closures of non-empty small sets are again models in a rosy theory T .

We review some notions and facts from [3],[4] and [8]. First we recall the definitions of simplices and the corresponding homology groups introduced in [3],[4]. **Throughout we work with a large saturated model $\mathcal{C}(=\mathcal{C}^{eq})$ whose theory $T(=T^{eq})$ is rosy with the thorn-independence relation $\perp$ on the small sets of $\mathcal{C}$. For a small $A \subset \mathcal{C}$, we denote the algebraic closure and definable closure in the home sort as $\text{acl}_{\mathcal{C}}(A)$ and $\text{dcl}_{\mathcal{C}}(A)$, and in the imaginary sort as $\text{acl}_{\mathcal{C}}^{eq}(A)$ and $\text{dcl}_{\mathcal{C}}^{eq}(A)$. If there is no risk of confusion, we shall write $\text{acl}(A)$ and $\text{dcl}(A)$ in both cases. We fix a small algebraically closed set $A = \text{acl}(A)$ and $p(x) \in S(A)$ (with possibly infinite x). When we say T is simple, we consider $T = T^{heq}$.**

Let $\mathcal{S}_A$ denote the category, where

- (1) The objects are small subsets of $\mathcal{C}$ containing A , and
- (2) The morphisms are elementary maps which fix A pointwise.

And for a finite $s \subset \omega$, the power set of s , $\mathcal{P}(s)$ forms the category as an ordered set :

- (1) $\text{Ob}(\mathcal{P}(s)) = \mathcal{P}(s)$, and
- (2) For $u, v \in \mathcal{P}(s)$, $\text{Mor}(u, v) = \{\iota_{u,v}\}$, where $\iota_{u,v}$ is the single inclusion map for $u \subseteq v$, or $= \emptyset$ otherwise.

For a functor $f : \mathcal{P}(s) \rightarrow \mathcal{C}_A$ and $u \subseteq v \in \mathcal{P}(s)$, we write $f_v^u := f(\iota_{u,v}) \in \text{Mor}(f(u), f(v))$ and $f_v^u(u) := f_v^u(f(u)) \subseteq f(v)$.

Definition 1.1. A functor $f : \mathcal{P}(s) \rightarrow \mathcal{S}_A$ for some finite $s \subset \omega$ is said to be a *closed independent (regular) n -simplex* in p if

- (1) $|s| = n + 1$
- (2) $f(\emptyset) \supseteq A$; and for $i \in s$, $f(\{i\})$ is of the form $\text{acl}(Ca)$ where $a \models p$ is independent with $C = f_{\{i\}}^{\emptyset}(\emptyset)$ over A .
- (3) For all non-empty $u \in \mathcal{P}(s)$, we have

$$f(u) = \text{acl}(A \cup \bigcup_{i \in u} f_{\{i\}}^{\{i\}}(\{i\}));$$

and $\{f_u^{\{i\}}(\{i\}) \mid i \in u\}$ is independent over $f_u^\emptyset(\emptyset)$.

We say f is *over* A if $f(\emptyset) = A$ (so for any $u \subset s$, $f_u^\emptyset(\emptyset) = A$). We shall call a closed independent n -simplex simply by an n -*simplex*. The set s is called the *support* of f , denoted by $\text{supp}(f)$.

In this paper, we only consider simplices over A . **We fix an enumeration of $\text{acl}(aA)$ for each $a \in \mathcal{C}^{|x|}$ such that $a, b \in \mathcal{C}^{|x|}$, $a \equiv_A b$ if and only if $\text{acl}(aA) \equiv \text{acl}(bA)$ because in [7], there is a counterexample which fails generalized amalgamation properties without fixing enumeration of bounded closed set.**

Definition 1.2. Let $S_n(p; A)$ denote the collection of all n -simplices over A in p and $C_n(p; A)$ the free abelian group generated by n -simplices in $S_n(p; A)$; its elements are called n -*chains* over A in p .

A non-zero n -chain c is uniquely written (up to permutation of i 's) as $c = \sum_{1 \leq i \leq k} n_i f_i$, where n_i is a non-zero integer and $f_1, \dots, f_k$ are distinct n -simplices. We call $|c| := |n_1| + \dots + |n_k|$ the *length* of the chain c , and define the *support* of c as the union of $\text{supp}(f_i)$'s.

We use $a, b, c, \dots, f, g, h, \dots, \alpha, \beta, \dots$ to denote simplices and chains. Now we define the boundary operators and using the boundary operators we will define homology groups.

Definition 1.3. Let $n \geq 1$ and $0 \leq i \leq n$. The i -th *boundary operator* $\partial_n^i : C_n(p; A) \rightarrow C_{n-1}(p; A)$ is defined so that if f is an n -simplex with domain $\mathcal{P}(s)$ with $s = \{s_0 < \dots < s_n\}$, then

$$\partial_n^i(f) = f \upharpoonright \mathcal{P}(s \setminus \{s_i\})$$

and extended linearly to all n -chains in $C_n(p)$.

The *boundary map* $\partial_n : C_n(p; A) \rightarrow C_{n-1}(p; A)$ is defined by the rule

$$\partial_n(c) = \sum_{0 \leq i \leq n} (-1)^i \partial_n^i(c).$$

We write ∂^i and ∂ for ∂_n^i and ∂_n , respectively, if n is clear from context.

Definition 1.4. The kernel of ∂_n is denoted $Z_n(p; A)$, and its elements are called (n) -*cycles* over A . The image of ∂_{n+1} in $C_n(p; A)$ is denoted by $B_n(p; A)$ and its elements are called (n) -*boundaries* over A .

Since $\partial_n \circ \partial_{n+1} = 0$, $B_n(p; A) \subseteq Z_n(p; A)$ and we can define simplicial homology groups in p .

Definition 1.5. The n -th (*simplicial*) *homology group* over A in p is

$$H_n(p; A) := Z_n(p; A) / B_n(p; A).$$

In [4], original simplicial homology groups in p were defined using not only simplices over A in p but also other simplices in p . But in this paper, we consider the homology groups over A in p .

Notation 1.6. We shall abbreviate $S_n(p; A), C_n(p; A), \dots$ as $S_n(p), C_n(p), \dots$ and we shall also abbreviate $H_n(p; A)$ simply as $H_n(p)$.

Definition 1.7. For $n \geq 1$, an n -chain c is called an n -shell if it is in the form

$$c = \pm \sum_{0 \leq i \leq n+1} (-1)^i f_i,$$

where $f_0, \dots, f_{n+1}$ are n -simplices such that whenever $0 \leq i < j \leq n+1$, we have $\partial^i f_j = \partial^{j-1} f_i$. Specially, a 1-shell c is of the form

$$c = f_0 - f_1 + f_2.$$

Remark 1.8. The boundary of an 2-simplex is a 1-shell, and the boundary of any 1-shell is 0.

Definition 1.9. For $n \geq 0$, we say p has $(n+2)$ -amalgamation if any n -shell in p is the boundary of some $(n+1)$ -simplex in p , and p has $(n+2)$ -complete amalgamation (or simply $(n+2)$ -CA) if p has k -amalgamation for every $2 \leq k \leq n+2$. By extension axiom of the independence relation, whenever $f : \mathcal{P}(s) \rightarrow \mathcal{C}_A, g : \mathcal{P}(t) \rightarrow \mathcal{C}_A \in S(p)$ and $f \upharpoonright \mathcal{P}(s \cap t) = g \upharpoonright \mathcal{P}(s \cap t)$, then f and g can be extended to a simplex $h : \mathcal{P}(s \cup t) \rightarrow \mathcal{C}_A$ in p . This property is called *strong 2-amalgamation*.

The following fact shows why the notion of shells is important.

Fact 1.10. [3],[4] *If p has $(n+1)$ -CA for some $n \geq 1$, then*

$H_n(p) = \{[c] : c \text{ is an } n\text{-shell over } A \text{ with } \text{supp}(c) = \{0, \dots, n+1\}\}.$
Thus the first homology group in p is generated by 1-shells in p with its support $\{0, 1, 2\}$.

So, we have that $H_1(p)$ is trivial iff any 1-shell of its support $\{0, 1, 2\}$ in p is the boundary of some 2-chain in p . Therefore, if T is simple, due to 3-amalgamation $H_1(p)$ is trivial. The following shows that the same result holds in any rosy theory.

Fact 1.11. [8] *Suppose that p is any Lascar strong type in a rosy theory. Then $H_1(p) = 0$.*

We introduce a notion of type homologies in [4]. We call types with possibly infinite sets of variables $*$ -types. We fix a set $\mathcal{V}$ of variables which is large enough so that all variables in $*$ -types come from the set $\mathcal{V}$ and $|\mathcal{C}| > 2^{|\mathcal{V}|}$. For any $X \subset \mathcal{V}$, any injective function $\sigma : X \rightarrow \mathcal{V}$,

and any $*$ -type $p(\bar{x})$ with $\bar{x} \subset X$, we let $\sigma_*p := \{\phi(\sigma(\bar{x})) : \phi(\bar{x}) \in p\}$. For $A = \text{acl}(A)$, let $\mathcal{T}_A$ be the category, where

- (1) The objects of $\mathcal{T}_A$ are all the complete $*$ -types in T over A , including a single distinguished type $p_\emptyset$ with no free variables;
- (2) $\text{Mor}_{\mathcal{T}_A}(p(\bar{x}), q(\bar{y}))$ is the set of all injective maps $\sigma : \bar{x} \rightarrow \bar{y}$ such that $\sigma_*p \subset q$.

Definition 1.12. Let $A = \text{acl}(A)$ and $p \in S(A)$. A *closed independent type- n -simplex in p* is a functor $f : \mathcal{P}(s) \rightarrow \mathcal{T}_A$ for $s \subset \omega$ such that

- (1) $|s| = n + 1$.
- (2) Let $w \subset s$ and $u, v \subset w$. Set $f_w^u := f(\iota_{u,w})$. Write $\bar{x}_w$ as the variable set of $f(w)$. Then whenever $\bar{a}$ realizes the type $f(w)$ and $\bar{a}_u, \bar{a}_v$, and $\bar{a}_{u \cap v}$ denote subtuples corresponding to the variable sets $f_w^u(\bar{x}_u)$, $f_w^v(\bar{x}_v)$, and $f_w^{u \cap v}(\bar{x}_{u \cap v})$, then

$$\bar{a}_u \downarrow_{A \cup \bar{a}_{u \cap v}} \bar{a}_v.$$

- (3) For all non-empty $u \subset s$ and any $\bar{a}$ realizing $f(u)$, we have $\bar{a} = \text{acl}(A \cup \bigcup_{i \in u} \bar{a}_{\{i\}})$.
- (4) For $i \in s$, $f(\{i\})$ is the complete $*$ -type of $\text{acl}(AC \cup \{b\})$ over A , where C is some realization of $f(\emptyset)$ and b is some realization of a nonforking extension of p to AC .

We say f is *over A* if $f(\emptyset) = A$.

Using closed independent type-functors in p we define the n -th type homology groups over A in p , denoted by $H_n^t(p; A)$. We shall write $H_n^t(p; A)$ as $H_n^t(p)$. Then for each n , the n -homology groups $H_n(p; A)$ and $H_n^t(p; A)$ are non-canonically isomorphic, which is depending on the choice of enumerations of each $*$ -types of closed independent simplices in p .

We see a notion of chain-walk notion, which is motivated from directed walk in graph theory, in [8][9]. The chain-walk was used to reduce a given a 2-chain of 1-shell boundary to one of simple form of 2-chain having the same 1-shell boundary and it is useful to compute the first homology group of a strong type. Two fundamental operations were used in reducing 2-chains to the forms of chain-walks : *crossing* and *renaming support* operations. We refer the reader to [8] for the definitions of crossing and renaming support operations and to [8][9] for the detail of classification of 2-chains. In [9], one defined the chain-walk using notion of direct walk in graph theory, here we give the definition of chain-walk in terms of simplices.

Definition 1.13. Let α be a 2-chain having the boundary $f_{12} - f_{02} + f_{01}$.

A subchain $\beta = \sum_{i=0}^m \epsilon_i b_i$ of α (where $\epsilon_i = \pm 1$ and b_i is a 2-simplex, for each i) is called a *chain-walk in α from f_{01} to $-f_{02}$* if

- (1) there are non-zero numbers $k_0, \dots, k_{m+1}$ (not necessarily distinct) such that $k_0 = 1$, $k_{m+1} = 2$, and for $i \leq m$, $\text{supp}(b_i) = \{k_i, k_{i+1}, 0\}$;
- (2) $(\partial \epsilon_0 b_0)^{0,1} = f_{01}$, $(\partial \epsilon_m b_m)^{0,2} = -f_{02}$; and
- (3) for $0 \leq i < m$,

$$(\partial \epsilon_i b_i)^{0,k_{i+1}} + (\partial \epsilon_{i+1} b_{i+1})^{0,k_{i+1}} = 0.$$

Any 2-chain having a 1-shell boundary is reduced to a chain-walk 2-chain having the same boundary of the support $\{0, 1, 2\}$.

Fact 1.14. [8][9] *Applying crossing and renaming support operation to a 2-chain α with the 1-shell boundary $f_{12} - f_{02} + f_{01}$, it is reduced to a 2-chain $\alpha' = \sum_{i=0}^{2n} (-1)^i a_i$ with $|\alpha'| \leq |\alpha|$, which itself is a chain-walk from f_{01} to $-f_{02}$, and $\text{supp}(\alpha') = \{0, 1, 2\}$.*

2. LASCAR GROUPS AND THE FIRST HOMOLOGY GROUPS

In this section we show that there is a canonical epimorphism from the Lascar group $\text{Gal}_L(\mathcal{C}; A)$ over A into the first homology group $H_1(p)$ in p .

Let $f: \mathcal{P}(s) \rightarrow \mathcal{C}_A$ be a n -simplex in p . For $u \subset s$ with $u = \{i_0 < \dots < i_k\}$, we shall write $f(u) = [a_0 \dots a_k]_u$, where $a_j \models p$, $f(u) = \text{acl}(A, a_0 \dots a_k)$, and $\text{acl}(a_j A) = f_u^{\{i_j\}}(\{i_j\})$, or we write $f(u) \equiv [a_0 \dots a_k]_u$ by changing '=' to ' $\equiv$ '. In both cases, $\{a_0, \dots, a_k\}$ is independent over A .

2.1. Representations of 1-shells. Given two 1-shells $s_k = f_{01}^k + f_{12}^k - f_{02}^k$ with $k = 0, 1$, if $f_{ij}^0(\{i, j\}) \equiv_A f_{ij}^1(\{i, j\})$ for $0 \leq i < j \leq 2$, then the homology classes of s_0 and s_1 are same in $H_1(p)$ since $H_1^t(p) \cong H_1(p)$. From this, we introduce a notion of a representation of a 1-shell and we describe the first homology group in p using this notion.

Definition/Remark 2.1. Let $s = f_{01} + f_{12} - f_{02}$ be a 1-shell such that $\text{supp}(f_{ij}) = \{i, j\}$ for $0 \leq i < j \leq 2$. There is a quadruple (a_0, a_1, a_2, a_3) in $p(\mathcal{C})^4$ such that $f_{01}(\{0, 1\}) \equiv [a_0 a_1]_{\{0,1\}}$, $f_{12}(\{1, 2\}) \equiv [a_1 a_2]_{\{1,2\}}$, and $f_{02}(\{0, 2\}) \equiv [a_3 a_2]_{\{0,2\}}$. We call this quadruple a *representation of s* .

Note that a representation for a 1-shell need not be unique and it is possible that the same quadruple represents different 1-shells even

though they have the same support because given $a, b \models p$, the enumeration of $\text{acl}(aA)$ is fixed but the enumeration of $\text{acl}(abA)$ is not fixed.

Definition 2.2. Let s be a 1-shell and (a, b, c, a') be a representation of s . We call a an *initial point*, a' a *terminal point*, (a, a') an *endpoint pair* of this representation.

In the next theorem, we'll see that the endpoint pairs of representations determine the classes of 1-shells in $H_1(p)$, and the group structure of $H_1(p)$ can be described by endpoint pairs.

Theorem 2.1. Let s_0 and s_1 be 1-shells of the support $\{0, 1, 2\}$. Suppose they have some representations of a same endpoint pair. Then $s_0 - s_1$ is a boundary of a 2-chain, that is, they are in the same homology class in $H_1(p)$

Proof. We assume that $A = \emptyset$. Consider two 1-shells $s_k = f_{12}^k - f_{02}^k + f_{01}^k$ for $k = 0, 1$. Suppose s_0 and s_1 have representations (a, b_0, c_0, a') and (a, b_1, c_1, a') respectively. Take two independent elements $b, c \models p$ such that $bc \perp ab_0b_1c_0c_1a'$ and consider a 1-shell s of its support $\{0, 3, 4\}$ which is represented by (a, b, c, a') . Then there is a 2-chain $\alpha = (a_{01}^0 + a_{12}^0 - b^0 - a_{02}^0) - (a_{01}^1 + a_{12}^1 - b^1 - a_{02}^1)$ where for each $k = 0, 1$ and $0 \leq i < j \leq 2$, a_{ij}^k and b^k are 2-simplices satisfying the followings :

- (1) If $j - i = 1$, $\text{supp}(a_{ij}^k) = \{i, j, 3\}$, otherwise, $\text{supp}(a_{02}^k) = \{0, 2, 4\}$. And $\text{supp}(b^k) = \{2, 3, 4\}$;
- (2) $a_{ij}^k \upharpoonright \mathcal{P}(\{i, j\}) = f_{ij}^k$; and
- (3) $a_{01}^k(\{0, 1, 3\}) = [ab_kb]_{\{0,1,3\}}$, $a_{12}^k(\{1, 2, 3\}) = [b_kc_kb]_{\{1,2,3\}}$, $a_{02}^k(\{0, 2, 4\}) = [a'c_kc]_{\{0,2,4\}}$; and $b^k(\{2, 3, 4\}) = [c_kbc]_{\{2,3,4\}}$.

and $\partial(a_{01}^k + a_{12}^k - b^k - a_{02}^k) = s_k - s$. So $\partial\alpha = (s_0 - s) - (s_1 - s) = s_0 - s_1$, and s_0 and s_1 are in the same homology class. $\square$

Theorem 2.2. Let s_0 and s_1 be 1-shells of a support $\{0, 1, 2\}$, and let $a, a', a'' \models p$ be such that (a, a') and (a', a'') are endpoint pairs of representations of s_0 and s_1 respectively. Then there is a 1-shell s of a support $\{0, 1, 2\}$ having an endpoint of representation (a, a'') and $[s] = [s_0] + [s_1]$ in $H_1(p)$.

Proof. Assume $A = \emptyset$. Consider two 1-shells of a support $\{0, 1, 2\}$, $s_0 = f_{01}^0 + f_{12}^0 - f_{02}^0$, $s_1 = f_{01}^1 + f_{12}^1 - f_{02}^1$. Suppose there are representations of s_0, s_1 such that the terminal point of one of s_0 and the initial point of one of s_1 are same. Let $a' \models p$ be the common element and $a, a'' \models p$ be elements so that (a, a') and (a', a'') are endpoint pairs of s_0 and s_1 respectively. Let $b_0, b_1, c_0, c_1 \models p$ be

elements such that two quadruples (a, b_0, c_0, a') and (a', b_1, c_1, a'') are representations of s_0 and s_1 respectively. Consider two independent elements $d, e \models p$ with $de \perp aa'a''b_0b_1c_0c_1$. Then there is a 2-chain $\alpha = (a_{01}^0 + a_{12}^0 - a_{02}^0) - b + (a_{01}^1 + a_{12}^1 - a_{02}^1)$, where for $k = 0, 1$ and $0 \leq i < j \leq 2$, a_{ij}^k and b are 2-simplices satisfying the followings :

- (1) $\text{supp}(a_{ij}^k) = \{i, j, 3+k\}$ and $\text{supp}(b) = \{0, 3, 4\}$;
- (2) $a_{ij}^k \upharpoonright \mathcal{P}(\{i, j\}) = f_{ij}^k$;
- (3) $a_{01}^0(\{0, 1, 3\}) = [a, b_0, d]_{\{0,1,3\}}$, $a_{12}^0(\{1, 2, 3\}) = [b_0, c_0, d]_{\{0,2,3\}}$,
 $a_{02}^0(\{0, 2, 3\}) = [a', c_0, d]_{\{0,2,3\}}$, $a_{01}^1(\{0, 1, 4\}) = [a', b_1, e]_{\{0,1,4\}}$,
 $a_{12}^1(\{1, 2, 4\}) = [b_1, c_1, e]_{\{1,2,4\}}$, $a_{02}^1(\{0, 2, 4\}) = [a'', c_1, e]_{\{0,2,4\}}$,
 and $b(\{0, 3, 4\}) = [a', d, e]_{\{0,3,4\}}$.

and $\partial(a) = s_0 + s_1 - s'$, where $s' = -a_{01}^0 \upharpoonright \mathcal{P}(\{0, 3\}) + b \upharpoonright \mathcal{P}(\{3, 4\}) + a_{12}^1 \upharpoonright \mathcal{P}(\{0, 4\})$ is a 1-shell of a support $\{0, 3, 4\}$. Using the proof of Theorem 2.1, we get a 1-shell s of a support $\{0, 1, 2\}$ having an endpoint (a, a'') and $[s] = [s']$ in $H_1(p)$. Thus, there is a 2-chain α' having a 1-chain $s_0 + s_1 - s$ as its boundary and so $[s] = [s_0] + [s_1]$. $\square$

Next we consider an action of $\text{Aut}_A(\mathcal{C})$ on each $C_n(p)$ and this action induces an action of $\text{Aut}_A(\mathcal{C})$ on $H_n(p)$. From the theorem 2.1, this action becomes trivial on $H_n(p)$. But this triviality is very crucial in finding a connection between the Lascar group over A and the first homology group in p .

Definition/Remark 2.3. We define an action of $\text{Aut}_A(\mathcal{C})$ on each $C_n(p)$. Let $\sigma \in \text{Aut}(\mathcal{C})$. For a n -chain $c = \sum_{i=0}^k n_i f_i$, we define

$$\sigma(c) := \sum_{i=0}^k n_i \sigma(f_i),$$

where for a n -simplex $f : \mathcal{P}(s) \rightarrow C_A$ with $s = \{s_0 < s_1 < \dots < s_n\}$, a n -simplex $\sigma(f)$ is defined as follows :

- (1) $\sigma(f)(u) := \sigma(f(u))$ for each $u \subset s$; and
- (2) $\sigma(f)(\iota_{u,v}) := \sigma \circ f(\iota_{u,v}) \circ \sigma^{-1}$ for each inclusion map $\iota_{u,v}$.

Furthermore, this action commutes with ∂ , i.e.,

$$\partial(\sigma(c)) = \sigma(\partial(c)).$$

So this action induces an action of $\text{Aut}_A(\mathcal{C})$ on $H_1(p)$ as follows : for each $[s] \in H_1(p)$, $\sigma([s]) := [\sigma(s)]$.

Note 2.4. Let s be a 1-shell in p and let (a, b) be an endpoint pair of s . For each $\sigma \in \text{Aut}_A(\mathcal{C})$, $(\sigma(a), \sigma(b))$ is an endpoint pair of $\sigma(s)$.

Since the n -th type-homology group and the n -th homology group in p are isomorphic, the action of $\text{Aut}_A(\mathcal{C})$ on $H_1(p)$ is trivial.

Corollary 2.5. *Let s be a 1-shell and let $\sigma \in \text{Aut}_A(\mathcal{C})$. Then there is a 2-chain α having the boundary of $s - \sigma(s)$.*

We denote the ordered bracket $[a, b]$ for the class of 1-shell s in $H_1(p)$ which has an endpoint pair (a, b) for $a, b \models p$. By Theorem 2.1, this bracket notion is well-defined. We can summarize Theorems 2.1, 2.2, and Corollary 2.5 as follows : For $a, b, c \in p(\mathcal{C})$ and $\sigma \in \text{Aut}_A(\mathcal{C})$, in $H_1(p)$,

- (1) $[a, b] + [b, c] = [a, c]$;
- (2) $[a, a]$ is the identity element;
- (3) $-[a, b] = [b, a]$; and
- (4) $\sigma([a, b]) = [\sigma(a), \sigma(b)] = [a, b]$.

2.2. Lascar group and the first homology groups. Here, using the ordered bracket notion of endpoint pairs, we define a map ψ_a from the automorphism group over A into the first homology group in p for each $a \models p$. This map is proven to be a surjective homomorphism (or epimorphism) and this map does not depend on the choice of $a \models p$. Thus we get a canonical epimorphism from $\text{Aut}_A(\mathcal{C})$ into $H_1(p)$ and we study about its kernel.

For each $a \models p$, we define a map ψ_a from $\text{Aut}_A(\mathcal{C})$ to $H_1(p)$ by sending σ into $[a, \sigma(a)]$.

Theorem 2.3. (1) *Each ψ_a is a epimorphism;*
 (2) *For $a, b \models p$, $\psi_a = \psi_b$. So we get a canonical map ψ from $\text{Aut}_A(\mathcal{C})$ into $H_1(p)$.*

Proof. (1) Fix $a \models p$. At first, surjectivity of ψ_a comes from the fact that for $b \models p$, there is a $\sigma \in \text{Aut}_A(\mathcal{C})$ such that $\sigma(a) = b$. It is enough to show that ψ_a is a homomorphism. For $\sigma, \tau \in \text{Aut}_A(\mathcal{C})$,

$$\begin{aligned}
 \psi(\sigma\tau) &= [a, \sigma\tau(a)] \\
 &= [a, \sigma(a)] + [\sigma(a), \sigma\tau(a)] \\
 &= [a, \sigma(a)] + \sigma[a, \tau(a)] \\
 &= [a, \sigma(a)] + [a, \tau(a)] \\
 &= \psi_a(\sigma) + \psi_a(\tau).
 \end{aligned}$$

So ψ_a is a homomorphism.

(2) Choose $a, b \models p$. Then there is $\tau \in \text{Aut}_A(\mathcal{C})$ such that $b = \tau(a)$.
For $\sigma \in \text{Aut}(\mathcal{C})$,

$$\begin{aligned}\psi_b(\sigma) &= \psi_a(\tau^{-1}\sigma\tau) \\ &= \psi_a(\tau^{-1}) + \psi_a(\sigma) + \psi_a(\tau) \\ &= \psi_a(\sigma).\end{aligned}$$

Thus $\psi_a = \psi_b$ and we get a canonical epimorphism $\psi(= \psi_a) : \text{Aut}_A(\mathcal{C}) \rightarrow H_1(p)$ for some $a \models p$. $\square$

So $H_1(p)$ is isomorphic to $\text{Aut}_A(\mathcal{C}) / \text{Ker}(\psi)$ and it is need to understand the kernel of ψ . In [8], it was shown that if p is a Lascar strong type, then the first homology group is zero. This fact can be restated using endpoint notion as follows :

Fact 2.6. [8] *Let $a, b \models p$ be such that $a \equiv_A^L b$. Then any 1-shell having endpoint pair (a, b) is a boundary of a 2-chain, i.e., $[a, b] = 0$ in $H_1(p)$.*

Definition 2.7. (1) Let $\text{Aut}_B(\mathcal{C})$ be the set of elements $\sigma \in \text{Aut}(\mathcal{C})$ fixing B pointwise.
(2) For a group G , the commutator of G is the subgroup of G generated by $\{ghg^{-1}h^{-1} \mid g, h \in G\}$, denoted by $[G, G]$ and this is the smallest normal subgroup between normal subgroups N of G making G/N abelian.

Theorem 2.4. *Let N be the normal subgroup of $\text{Aut}_A(\mathcal{C})$ generated by automorphisms in $\text{Autf}_A(\mathcal{C})$, or in $\text{Aut}_{\text{acl}(Aa)}(\mathcal{C})$ for some $a \models p$, let $G = \text{Aut}_A(\mathcal{C})/N$, and consider the canonical quotient map $\Psi' : \text{Aut}_A(\mathcal{C}) \rightarrow G$. Let $\psi : \text{Aut}_A(\mathcal{C}) \rightarrow H_1(p)$ be the canonical map. Then the kernel of ψ contains the followings :*

- (1) $\text{Aut}_{\text{acl}(Aa)}(\mathcal{C})$ for each $a \models p$;
- (2) $\text{Autf}_A(\mathcal{C})$; and
- (3) $(\Psi')^{-1}([G, G])$.

Specially, from the second one, ψ induces a canonical epimorphism Ψ from $\text{Gal}_L(\mathcal{C}; A)$ into $H_1(p)$.

Proof. (1) For any $a \models p$, $[a, a]$ is the identity element in $H_1(p)$, and since we fix an enumeration of $\text{acl}(aA)$, $\text{Aut}_{\text{acl}(Aa)}(\mathcal{C})$ is contained the kernel of ψ .

(2) It comes from Fact 2.6

(3) It comes from the fact that $H_1(p)$ is always abelian. $\square$

We define an A -invariant equivalence relation on $p(\mathcal{C})^2$, and using this equivalence relation we describe the kernel of ψ . By Fact 1.14, we can describe 1-shells which are boundary of 2-chains as follows :

Theorem 2.5. *A 1-shell s is a boundary of 2-chain if and only if there is a representation (a, b, c, a') such that for some n there is a finite sequence $(d_i)_{0 \leq i \leq 2n+1}$ of elements in $p(\mathcal{C})$ satisfying the following conditions :*

- (1) $d_0 = a, d_{2n+1} = c$ and $d_{2i_0} = a'$ for some $0 < i_0 \leq n$;
- (2) $\{d_{2i}, d_{2i+1}, b\}$ is independent for each $0 \leq i \leq n$; and
- (3) There is a bijection m from $\{0, 2, \dots, 2n\} \setminus \{2i_0\}$ to $\{1, 3, \dots, 2n-1\}$ such that $d_{2i}d_{2i+1} \equiv d_{m(2i)+1}d_{m(2i)}$ for $0 \leq i \neq i_0 \leq n$, and $d_{2i_0}d_{2i_0+1} \equiv a'c$.

Proof. Let $s = f_{01} + f_{12} - f_{02}$ be a 1-shell and let $\alpha = \sum_{i=0}^{2n} (-1)^i a_i$ be a 2-chain with $\text{supp}(\alpha') = \{0, 1, 2\}$ and $\partial(\alpha) = s$, which is a chain-walk from f_{01} to $-f_{02}$. Then there are a representation (a, b, c, a') and a finite sequence $(d_i)_{0 \leq i \leq 2n+1}$ of elements in $p(\mathcal{C})$ satisfying the following conditions :

- (1) $d_0 = a, d_{2n+1} = c$ and $d_{2i_0} = a'$ for some $0 < i_0 \leq n$;
- (2) $\{d_{2i}, d_{2i+1}, b\}$ is independent for each $0 \leq i \leq n$; and
- (3) There is a bijection m from $\{0, 2, \dots, 2n\} \setminus \{2i_0\}$ to $\{1, 3, \dots, 2n-1\}$ such that $d_{2i}d_{2i+1} \equiv d_{m(2i)+1}d_{m(2i)}$ for $0 \leq i \neq i_0 \leq n$, and $d_{2i_0}d_{2i_0+1} \equiv a'c$.

□

Now we define an A -invariant equivalence relation on $p(\mathcal{C})$ representing the kernel of ψ . For $m \geq 1$, define a partial type $p^{\odot m}(x_1, \dots, x_m)$ over A as

$$\bigwedge_{i \leq m} p(x_i) \wedge \bigwedge_{j < m} x_j \perp x_{j+1}.$$

Next define a relation $\sim$ on $p^{\odot 4}(\mathcal{C})$ as follows : for $(a_i, b_i, c_i, a'_i) \in p^{\odot 4}(\mathcal{C})$ and $i = 0, 1$, $(a_0, b_0, c_0, a'_0) \sim (a_1, b_1, c_1, a'_1)$ if $a_0b_0 \equiv_A a_1b_1$, $b_0c_0 \equiv_A b_1c_1$, and $a'_0c_0 \equiv_A a'_1c_1$. Note that for (a_0, b_0, c_0, a'_0) and (a_1, b_1, c_1, a'_1) in $p^{\odot 4}(\mathcal{C})$, $(a_0, b_0, c_0, a'_0) \sim (a_1, b_1, c_1, a'_1)$ if and only if both quadruples represent a same 1-shell. This relation is an A -type-definable equivalence relation. And for each $n \geq 0$, the (1), (2) and (3) conditions of (a, b, c, a') and $(d_0, d_1, \dots, d_{2n+1})$ in Theorem 2.5 is A -type-definable.

We define a partial type $F_n(x, y, z, w; v_0, v_1, \dots, v_{2n+1})$ as

$$\begin{aligned}
 & \bigwedge_{0 \leq i \leq 2n+1} p(v_i) \wedge v_0 = x \wedge v_{2n+1} = w \\
 & \wedge \bigwedge_{0 \leq j \leq n} \{v_{2j}, v_{2j+1}, z\} \text{ is independent over } A \\
 & \wedge \bigvee_{0 \leq i_0 \leq n} [d_{2i_0} = y] \\
 & \wedge \bigvee_m \left(\bigwedge_{0 \leq i \neq i_0 \leq n} v_{2i} v_{2i+1} \equiv_A v_{m(2i)+1} v_{m(2i)} \right) \\
 & \quad \text{for each bijection } m : \{0, 2, \dots, 2n\} \setminus \{2i_0\} \rightarrow \{1, 3, \dots, 2n-1\} \\
 & \wedge v_{2i_0} v_{2i_0+1} \equiv_A yw].
 \end{aligned}$$

At last, for each $n \geq 0$, we define a partial type $E'_n(x, y)$ over A as

$$\begin{aligned}
 & p(x) \wedge p(y) \\
 & \wedge \exists z w x' y' z' w' [(p^{\odot 4}(x, z, w, y) \wedge p^{\odot 4}(x', z', w', y') \wedge (x, z, w, y) \sim (x', z', w', y')) \\
 & \wedge \exists v_0 v_1 \dots v_{2n+1} F_n(x', y', z', w'; v_0, v_1, \dots, v_{2n+1})].
 \end{aligned}$$

The relation $E'_n(x, y)$ says that (x, y) is an endpoint pair of a 1-shell which is a boundary of 2-chain which is a chain-walk of length $2n+1$. Take $E_n(x, y) \equiv E'_n(x, y) \wedge E'_n(y, x)$. So, for each $n \geq 0$, E_n is A -type-definable symmetric relation. At last, define the binary relation $E(x, y)$ as

$$x = y \vee \bigvee_{n \geq 0} E_n(x, y).$$

This relation is A -invariant, reflexive, and symmetric. By Theorem 2.2, it is transitive and by Theorem 2.5 $E(a, b)$ if and only if $[a, b] = 0$ in $H_1(p)$ for $a, b \models p$. So this relation E is a desired A -invariant equivalence relation.

Next, we define a distance-like notion on $p(\mathcal{C})$ as follows : For $a, b \models p$,

$$d_E(a, b) := \begin{cases} \min\{n \mid E_n(a, b)\} & \text{if } E(a, b) \\ \infty & \text{otherwise.} \end{cases}$$

This distance-like notion is not necessary to satisfy triangle inequality, i.e., for $a, b, c \models p$, $d_E(a, b) \leq d_E(a, c) + d_E(c, b)$. But it does hold that for $a, b, c \models p$, $d_E(a, b) \leq d_E(a, c) + d_E(c, b) + 8$ since in the proof of Theorem 2.2 for two 1-shells s_0 and s_1 , there is a 1-shell s such that $s_0 + s_1 - s$ is a boundary of 2-chain of length of $15 (= 2 \times 8 - 1)$. So we can apply the results in [11] and we know that if E is not type-definable, then the cardinality of $H_1(p)$ is at least $2^{\aleph_0}$. And, in the Appendix A, we prove that for any bounded (type-)definable equivalence relation on a strong type, the possible cardinality of the set of equivalence classes in the strong type is one or at least $2^{\aleph_0}$. In [10], an invariant equivalence

relation E on a type-definable set X is called *orbital equivalence relation* if there is a subgroup Γ of $\text{Aut}(\mathcal{C})$ such that Γ preserves classes of E setwise and acts transitively on each class. By Theorem 2.3, $E(a, b)$ if and only if there is $\sigma \in \text{Ker}(\psi)$ such that $\sigma(a) = b$ for $a, b \models p$. So, our equivalence relation E is a orbital equivalence relation.

Theorem 2.6. (1) E is an orbital equivalence relation.
 (2) The cardinality of $H_1(p)$ is zero or $\geq 2^{\aleph_0}$.

Next section, we give two examples and in two examples, we compute two first homology groups, which are non trivial and their cardinalities are exactly $2^{\aleph_0}$.

3. EXAMPLES

In simple theories including stable theories, the first homology group of a strong type is always zero by 3-amalgamation. In [4], the first homology groups of strong types were computed for some cases and they were all zero, and they showed that in o-minimal theories, the first homology group of a strong 1-type is always trivial. Here, we give two examples of rosy theories having a non trivial first homology group in a strong type. They are the first cases to give a non trivial first homology group in a strong type. In [8], B. Kim, S. Kim, and the author considered the structures in [1], $\mathcal{M}_{1,n} = (M; S; g_{1/n})$ for each $n \in \mathbb{N} \setminus \{0\}$ where

- (1) M is a saturated circle;
- (2) $g_{1/n}$ is a rotation (clockwise) by $2\pi/n$ -radian; and
- (3) S is a ternary relation such that $S(a, b, c)$ holds if a, b, c are distinct and b comes before c going around the circle clockwise starting at a .

and it was shown that the unique strong 1-type p_n in $S_1(\emptyset)$ has the trivial first homology group for every n , which is actually a Lascar strong type. Here we consider two structure $\mathcal{M}_1 = (M; S; g_{1/n} : n \in \mathbb{N} \setminus \{0\})$ expanding the structures $\mathcal{M}_{1,n}$ by adding all rotation functions of $2\pi/n$ -radian for each $n \in \mathbb{N} \setminus \{0\}$ at the same time. When we write g_r for $r = m/n$ in $\mathbb{Q} \cap [0, 1)$, it means $g_{1/n}^m$, and $\mathcal{M}_2 = (M; U_{<r}, U_{=r} | r \in (0, 1/2] \cap \mathbb{Q})$, where $U_{<r}(x, y)$ says the smallest length between x and y along the arc is less than $2\pi r$, and $U_{=r}(x, y)$ says the smallest length between x and y along the arc is exactly equal to $2\pi r$.

3.1. Rosiness of $\text{Th}(\mathcal{M}_1)$ and $\text{Th}(\mathcal{M}_2)$. In this subsection, we mainly show that two theories of $\mathcal{M}_1$ and $\mathcal{M}_2$ are rosy. In [2], C. Ealy and A. Onshuus gave a sufficient condition for being a rosy theory.

Fact 3.1. *Any theory T which geometrically eliminates imaginaries and for which algebraic closure defines a pregeometry is rosy of thorn U -rank 1.*

For rosiness of $\text{Th}(\mathcal{M}_i)$ ($i = 1, 2$), we show that $\text{Th}(\mathcal{M}_1)$ has weak elimination of imaginaries and $\text{Th}(\mathcal{M}_2)$ has geometric elimination of imaginaries. In [12], B. Poizat defined for a theory T to have *weak elimination of imaginaries* if for every definable set has a smallest algebraically closed set which it is definable over. And T has a *geometric elimination of imaginaries*, a weaker notion of weak elimination of imaginaries, if for each imaginary $e \in \mathcal{M}^{eq} \models T$, there is a real tuple $\bar{a} \subset M$ such that $e \in \text{acl}^{eq}(\bar{a})$ and $\bar{a} \in \text{acl}^{eq}(e)$. We give a sufficient condition of weak elimination of imaginaries for $\aleph_0$ -categorical theory, used in [8] :

Theorem 3.1. *Let T be $\aleph_0$ -categorical and let $\mathcal{M} = (M, \dots)$ be a saturated model of T . Suppose that for all A , $\text{acl}(A) = \text{dcl}(A)$. Suppose for a subset X of M^1 , if X is $A_0 (= \text{acl}(A_0))$ -definable and $A_1 (= \text{acl}(A_1))$ -definable, then X is $B (= A_0 \cap A_1)$ -definable. Then for a subset Y of M^n , if Y is A_0 -definable and A_1 -definable, then Y is B -definable. Furthermore, in this case, T has weak elimination of imaginaries.*

Proof. Let $A_0 = \text{acl}(A_0)$, $A_1 = \text{acl}(A_1)$, and $B = A_0 \cap A_1$. We use induction on n . If $n = 1$, it holds by assumption. Let's show this holds for the case $n + 1$ with inductive hypothesis for the case n . Let $A_0 = \text{acl}(A_0)$, $A_1 = \text{acl}(A_1)$, and $B = A_0 \cap A_1$. We may assume A_0 and A_1 are finite, and so is B . Let $Y \subset M^{n+1}$ be A_i -definable, defined by formula $\phi_i(x_0, \dots, x_n; \bar{a}_i)$ for $\bar{a}_i \subset A_i$ respectively. Then for each $c \in M$, the fiber of Y over c , $Y_c := \{\bar{x} \in M^n \mid \phi_i(\bar{x}, c; \bar{a})\}$ is cB -definable by induction. By $\aleph_0$ -categoricity, there are only finitely many formulas over $\emptyset$ modulo T , and it easily follows that for each y , $\phi_i(x_0, \dots, x_{n-1}, y, \bar{a}_i)$ is B -definable. Thus Y is B -definable.

And since there is no infinite descending chain of algebraically closed sets generating by finitely many elements, it makes for any definable set to have a smallest algebraically closed set where it is definable. Thus T weakly eliminate imaginaries. $\square$

As a corollary of Theorem 3.1, we showed that for each $n \geq 2$, $\text{Th}(\mathcal{M}_{1,n})$ has weak elimination of imaginaries.

Fact 3.2. [8] *For each $n \geq 2$, $\text{Th}(\mathcal{M}_{1,n})$ weakly eliminates imaginaries.*

Next we will see that the theory of $\mathcal{M}_1$ has quantifier-elimination.

Definition 3.3. Let M be the underlying set of $\mathcal{M}_1$ and $\mathcal{M}_2$, so it is a saturated circle. For $A \subset M$, let $\text{cl}(A) := \{g_r(a) \mid a \in A, r \in \mathbb{Q} \cap [0, 1)\}$.

Later, we will see that $\text{cl}(A) = \text{dcl}_{\mathcal{M}_1}(A) = \text{acl}_{\mathcal{M}_1}(A)$ in the home sort of $\mathcal{M}_1$ and $\text{cl}(A) = \text{acl}_{\mathcal{M}_2}$ in the home sort of $\mathcal{M}_2$. It is also easy to see that $\text{cl}(A)$ is a substructure of $\mathcal{M}_1$ and $\mathcal{M}_2$.

Theorem 3.2. *The theory of $\mathcal{M}_1$ has quantifier-elimination.*

Proof. Take two small subset $A, B \subset M$ such that $A = \text{cl}(A)$ and $B = \text{cl}(B)$ in M . Take $a \in M \setminus A$. We will find $b \in M \setminus B$ such that the map $f \cup \{(a, b)\}$ is extended to an embedding from $\text{cl}(Aa)$ to $\text{cl}(Bb)$ in $\mathcal{M}_1$. Then, the quantifier-elimination of $\text{Th}(\mathcal{M}_1)$ comes from a standard argument. We divide A into two parts $A_0 := \{x \in A \mid S(a, x, g_{1/2}(a))\}$ and $A_1 := \{x \in A \mid S(g_{1/2}(a), x, a)\}$. Then B is also divided into two parts $B_0 = f(A_0)$ and $B_1 = f(A_1)$. Take arbitrary $b \in M$ such that for all $y_0 \in B_0$, $y_1 \in B_1$, $S(y_1, b, y_0)$. Then b is a desired element. $\square$

Theorem 3.3. *The theory of $\mathcal{M}_1$ weakly eliminate imaginaries, and is rosy of thorn U -rank 1.*

Proof. In the structure $\mathcal{M}_1$, there is no infinite descending chain of algebraical closure of finite sets by quantifier elimination. It is enough to show that if $X \subset M^n$ is $A_0 (= \text{acl}(A_0))$ - and $A_1 (= \text{acl}(A_1))$ -definable, then X is $A_0 \cap A_1 (= B)$ -definable. Then X has a smallest algebraically closed set defining X , and $\text{Th}(\mathcal{M}_1)$ has weak elimination of imaginaries.

Let $A_i = \text{acl}(A_i) = \text{cl}(A_i)$ for $i = 0, 1$ and let $B = A_0 \cap A_1$. Let $X \subset M^m$ be A_i -definable in $\mathcal{M}_1$. Then X is definable over A_i for $i = 0, 1$ in some reduct $\mathcal{M}_{1,n}$. Since $\mathcal{M}_{1,n}$ weakly eliminates imaginaries, X is definable over B in $\mathcal{M}_{1,n}$, defined by a formula $\psi(\bar{x}, \bar{b})$. Then by the same formula $\psi(\bar{x}, \bar{b})$, X is B -definable in $\mathcal{M}_1$.

By quantifier elimination, it is easily verified that the algebraic closure in $\mathcal{M}_1$ gives a trivial pregeometry. Thus by Fact 3.1, $\text{Th}(\mathcal{M}_1)$ is a rosy theory having thorn U -rank 1. $\square$

There is only one 1-strong type over empty set $p_0(x) = \{x = x\}$ in $\mathcal{M}_1$.

Next we show that $\text{Th}(\mathcal{M}_2)$ has geometric elimination of imaginaries. We consider an expansion of $\mathcal{M}_2$, $\mathcal{N}_2 = (M, U_{<r}, U_{=r}, g_{1/n}) \mid r \in (0, 1/2] \cap \mathbb{Q}, n > 0$ by adding rotation functions $g_{1/n}$, and the reducts of $\mathcal{N}_2$, $\mathcal{N}_{2,k} = (M, U_{<r}, U_{=r}, g_{1/k}) \mid r \in (0, 1/2]_k$, where $(0, 1/2]_k = \{i/k \mid 0 < i/k \leq 1/2, i \in \mathbb{N}\}$, for each $k \geq 1$.

Theorem 3.4. (1) *For each $k \geq 3$, $\text{Th}(\mathcal{N}_{2,k})$ has quantifier-elimination, and weak elimination of imaginaries, and it is ω -categorical.*
 (2) *$\text{Th}(\mathcal{N})$ has weak elimination of imaginaries.*

Proof. (1) Fix $k \geq 3$. Each binary relations $U_{<r}$ and $U_{=r}$ for $r \in (0, 1/2]_k$ are $\emptyset$ -definable in $\mathcal{M}_{1,k}$ by quantifier-free formulas. So, by

Fact 3.2, it is enough to show that the ternary relation $S(x, y, z)$ is $\emptyset$ -definable in $\mathcal{N}_{2,k}$ by a quantifier-free formula. Denote $g_{i/k}(x) < y < g_{(i+1)/k}(x)$ for the formula $U_{<1/k}(g_{i/k}(x), y) \wedge U_{<1/k}(g_{(i+1)/k}(x), y)$. Consider the following quantifier-free formula

$$\begin{aligned} S'_k(x, y, z) \equiv & \bigvee_{i=1}^{k-1} [z = g_{i/k}(x) \rightarrow (\bigvee_{j=1}^{i-1} y = g_{j/k}(x) \\ & \vee \bigvee_{j=0}^{i-1} g_{j/k}(x) < y < g_{(j+1)/k}(x))] \\ & \vee \bigvee_{i=0}^{k-1} [g_{i/k}(x) < z < g_{(i+1)/k}(x) \rightarrow \\ & (g_{i/k}(x) < y < g_{(i+1)/k}(x) \wedge g_{-1/k}(z) < y < z) \\ & \vee \bigvee_{j=0}^{i-1} g_{j/k}(x) < y < g_{(j+1)/k}(x)], \end{aligned}$$

and this formula defines $S(x, y, z)$ in $\mathcal{N}_{2,k}$.

(2) Consider the ternary relation S'_k for some $k \geq 3$. Then S'_k also defines the ternary relation S in $\mathcal{N}$. Thus, as the relation between $\mathcal{N}_{2,k}$ and $\mathcal{M}_{1,k}$, by Theorems 3.2 and 3.3, the theory of $\mathcal{N}$ has quantifier-elimination and weakly eliminate imaginaries. $\square$

Theorem 3.5. *The theory of $\mathcal{M}_2$ geometrically eliminate imaginaries.*

Proof. At first, we define some equivalence relations E_n on M^n , which are $\emptyset$ -definable in $\mathcal{M}_2$. Let $n \geq 1$. Define a formula

$$C_n(z_1, \dots, z_n) \equiv \bigwedge_{1 \leq i < j \leq n} z_i \neq z_j \wedge \bigwedge_{1 \leq i < j \leq n} \neg U_{<1/n}(z_i, z_j)$$

so that it defines the set $\{z, g_{1/n}(z), \dots, g_{(n-1)/n}(z)\}$. Next, we define the following formula

$$\begin{aligned} I_n(x, y; z_1, \dots, z_n) \equiv & C_n(\bar{z}) \\ & \wedge \left[\bigvee_{i=1}^n (y = z_i) \rightarrow (U_{<1/n}(z_i, x) \wedge U_{<1/n}(z_{i+1}, x)) \right] \\ & \vee \left[\bigvee_{i=1}^n (U_{<1/n}(z_i, x) \wedge U_{<1/n}(z_{i+1}, x)) \rightarrow \right. \\ & \quad \left. \exists y_2 \cdots y_n (C_n(y, y_2, \dots, y_n) \right. \\ & \quad \wedge (U_{<1/n}(z_{i+1}, y_2) \wedge U_{<1/n}(z_{i+2}, y_2)) \\ & \quad \left. \wedge (U_{<1/n}(y, x) \wedge U_{<1/n}(y_2, x)) \right] \end{aligned}$$

, where $z_{n+1} = z_1$ and $z_{n+2} = z_2$, which defines the sets $\{(a, b) \mid b < a < g_{1/n}(b)\}$ or $\{(a, b) \mid g_{-1/n}(b) < a < b\}$ up to $z_i = g_{(i-1)/n}(z_1)$ or $z_i = g_{-(i-1)/n}(z_1)$ respectively. At last, consider the following equivalence

relation E_n on M^n defined as follows

$$E_n(\bar{z}, \bar{z}') \equiv \forall xy(I_n(x, y; \bar{z}) \leftrightarrow I_n(x, y; \bar{z}')).$$

Then $|M^n/E_n| = 3$ and each classes represents one of the following tuples :

$$\begin{cases} (z, g_{1/n}(z), \dots, g_{(n-1)/n}(z)) \\ (z, g_{-1/n}(z), \dots, g_{-(n-1)/n}(z)) \\ \text{other wise} \end{cases}$$

Let $\mathcal{L}(\mathcal{M}_2) = \{U_{<r}, U_{=r}\}$ be the language of $\mathcal{M}_2$ and $\mathcal{L}(\mathcal{N}_2) = \mathcal{L}(\mathcal{M}_2) \cup \{g_{1/n}\}_{n \geq 1}$ be the language of $\mathcal{N}_2$. Expand M_2 to $M'_2 = (M, U_{<r}, U_{=r}, a_n, b_n, c_n)_{n \geq 1}$ by adding imaginary elements such that $\{a_n, b_n, c_n\} = M^n/E_n$, where $a_n = [z, g_{1/n}(z), \dots, g_{(n-1)/n}(z)]_{E_n}$, and $b_n = [z, g_{-1/n}(z), \dots, g_{-(n-1)/n}(z)]_{E_n}$, for each $n \geq 1$, and the language of $\mathcal{M}_2$, $\mathcal{L}(\mathcal{M}_2)$ is the union of $\mathcal{L}(\mathcal{M}_2)$ and $\{S_{E_n}, f_{E_n}\}_{n \geq 1}$, where S_{E_n} is interpreted as the sort for M^n/E_n and f_{E_n} is as the canonical function from M_n into S_{E_n} such that for $\bar{a}, \bar{b} \in M^n$, $f_{E_n}(\bar{a}) = f_{E_n}(\bar{b})$ if and only if $E_n(\bar{a}, \bar{b})$.

Claim 3.4. *Each function $g_{1/n}$ is definable in M'_2 .*

Proof. Two functions g_1 and $g_{1/2}$ are already definable in M_2 and so in M'_2 also. Let $n \geq 3$. For each $a \in M$, there is only one element a' in M such that $\exists x_3, \dots, x_n (C_n(a, a', x_3, \dots, x_n) \wedge f_{E_n}(a, a', x_3, \dots, x_n) = a_n)$, and thus $a' = g_{1/n}(a)$. Therefore, the graph of $g_{1/n}$ is defined by the formula $\exists x_3, \dots, x_n (C_n(x, y, x_3, \dots, x_n) \wedge f_{E_n}(x, y, x_3, \dots, x_n) = a_n)$. $\square$

Claim 3.5. $(\mathcal{M}'_2)^{eq} = \mathcal{N}_2^{eq}$.

Proof. For each $e \in (\mathcal{M}'_2)^{eq}$, since $M^n/E_n \subset \mathcal{M}_2^{eq}$, e is in $(\mathcal{M}_2^{eq})^{eq} = \mathcal{M}_2^{eq}$. By the way, $\mathcal{M}_2$ is a reduct of $\mathcal{N}_2$, and $e \in \mathcal{N}_2^{eq}$. Conversely, each $e' \in \mathcal{N}_2^{eq}$ is clearly in $(\mathcal{N}'_2)^{eq}$, where $\mathcal{N}'_2 = (M, U_{<r}, U_{=r}, g_{1/n}, a_n, b_n, c_n)$. By Claim 3.4, $(\mathcal{N}'_2)^{eq} = (\mathcal{M}'_2)^{eq}$ and $e' \in (\mathcal{M}'_2)^{eq}$. $\square$

Take $e \in \mathcal{M}_2^{eq}$ arbitrary. Since $\mathcal{M}_2$ is a reduct of $\mathcal{N}_2$, e is in $\mathcal{N}_2^{eq}$. By weak elimination of imaginaries of $\text{Th}(\mathcal{N}_2)$, there is a finite tuple $\bar{b} \subset M$ such that

$$e \in \text{dcl}_{\mathcal{N}_2}^{eq}(\bar{b}) \text{ and } \bar{b} \in \text{acl}_{\mathcal{N}_2}^{eq}(e).$$

By Claim 3.5,

$$e \in \text{dcl}_{\mathcal{M}'_2}^{eq}(\bar{b}) \text{ and } \bar{b} \in \text{acl}_{\mathcal{M}'_2}^{eq}(e).$$

Since $M^n/E_n \subset \text{acl}_{\mathcal{M}_2}^{eq}(\emptyset)$,

$$e \in \text{acl}_{\mathcal{M}_2}^{eq}(\bar{b}) \text{ and } \bar{b} \in \text{acl}_{\mathcal{M}_2}^{eq}(e).$$

Therefore, each imaginaries in $\mathcal{M}_2^{eq}$ is inter-algebraic with a finite tuple in the home sort in $\mathcal{M}_2$ and $\text{Th}(\mathcal{M}_2)$ has geometric elimination of imaginaries. $\square$

Now we show that $\text{Th}(\mathcal{M}_2)$ is a rosy theory having thorn- U rank 1.

Theorem 3.6. *The theory of $\mathcal{M}_2$ is a rosy theory of thorn- U rank 1.*

Proof. By Fact 3.1 and Theorem 3.5, it is enough to show that the algebraic closure in the home sort gives a trivial pregeometry in $\mathcal{M}_2$. For any $A \subset M$, it is clear that $\text{cl}(A) \subset \text{acl}_{\mathcal{M}_2}(A)$. By the way, from Theorem 3.4, $\text{acl}_{\mathcal{N}_2}(A) = \text{cl}(A)$. Since $\mathcal{M}_2$ is a reduct of $\mathcal{N}_2$, $\text{acl}_{\mathcal{M}_2}(A) \subset \text{acl}_{\mathcal{N}_2}(A)$, and thus $\text{cl}(A) = \text{acl}_{\mathcal{M}_2}(A)$. So, the algebraic closure in $\mathcal{M}_2$ gives a trivial pregeometry. $\square$

The theory of $\mathcal{M}_2$ does not eliminate quantifier but there is only one 1-type over $\text{acl}^{eq}(\emptyset) (\neq \emptyset)$, $q_0(x) = \{x = x\}$.

3.2. Computation of H_1 in $\mathcal{M}_1$. In the section 2, we see that the first homology groups are determined by end-point pairs of 1-shells. In $\mathcal{M}_1$, for a fixed $a \in M$, we observe that $S_1(a)$ looks like a circle with a rotation. From this observation, we compute the first homology group of p_0 in $\mathcal{M}_1$:

Theorem 3.7. *In $\mathcal{M}_1$, the first homology group of p_0 is isomorphic to $\mathbb{R}/\mathbb{Z}$.*

We start with defining a distance-like notion between two points on M . For a subset A in $\mathbb{R}$, we denote $A_{\mathbb{Q}}$ for $A \cap \mathbb{Q}$.

Definition 3.6. Let $a, b \in M$ be two elements. We define the S -distance of b from a , denoted by $\text{Sd}(a, b)$ as follows : For $r \in \mathbb{Q}$ and $s < t \in [0, 1)_{\mathbb{Q}}$,

- (1) $\text{Sd}(a, b) = r$ if $b = g_r(a)$;
- (2) $s < \text{Sd}(a, b) < 1$ if $\mathcal{M}_1 \models S(g_s(a), b, a)$;
- (3) $0 < \text{Sd}(a, b) < t$ if $\mathcal{M}_1 \models S(a, b, g_t(a))$; and
- (4) $s < \text{Sd}(a, b) < t$ if $\mathcal{M}_1 \models S(g_s(a), b, g_r(a))$.

For $r \in [0, 1) \setminus \mathbb{Q}$, we write $\text{Sd}(a, b) = r$ if for $s < t \in [0, 1)_{\mathbb{Q}}$ and $s < r < t$, $s < \text{Sd}(a, b) < t$. Let $r \in (0, 1)_{\mathbb{Q}}$. We write $\text{Sd}(a, b) = r - \epsilon$ if for $s \in (0, 1)_{\mathbb{Q}}$ with $s < r$, $s < \text{Sd}(a, b) < r$. We write $\text{Sd}(a, b) = r + \epsilon$ if for $t \in (0, 1)_{\mathbb{Q}}$ with $r < t$, $r < \text{Sd}(a, b) < t$. We write $\text{Sd}(a, b) = \epsilon$ if for all $s \in (0, 1)_{\mathbb{Q}}$, $0 < \text{Sd}(a, b) < s$. We write $\text{Sd}(a, b) = 1 - \epsilon$ if for all $s \in (0, 1)_{\mathbb{Q}}$, $s < \text{Sd}(a, b) < 1$.

For a subset $A \subset \mathbb{Q}$, we define $A^* := A \cup \{x \pm \epsilon \mid x \in A\}$. This S -distance has the values in $[0, 1) \cup [0, 1)_{\mathbb{Q}}^* \cup \{1 - \epsilon\}$. In Appendix B, using Dedekind

cut, we develop multivalued operations $+$, $\times$, $-$ to make $\mathbb{R} \cup \mathbb{Q}^*$ a ring-like structure. Now we extend the values of S -distance to $\mathbb{R} \cup \mathbb{Q}^*$. Since $g_k = \text{id}$ for all $k \in \mathbb{Z}$, we write $\text{Sd}(a, b) = r$ for $r \in \mathbb{R} \cup \mathbb{Q}^*$ if $\text{Sd}(a, b) = r'$ where r' is the unique number in $[0, 1) \cup [0, 1)_{\mathbb{Q}}^*$ such that $r \in r' +^* n$ for some $n \in \mathbb{Z}$. Then this values depends only on the type of (a, b) , that is, for $a_0, a_1, b_0, b_1 \in M$, if $a_0 b_0 \equiv a_1 b_1$, then $\text{Sd}(a_0, b_0) = \text{Sd}(a_1, b_1)$ of the values in $[0, 1) \cup [0, 1)_{\mathbb{Q}}^*$. Then the following fact is easily verified :

Fact 3.7. *Let $a, b, c \in M$.*

- (1) $\text{Sd}(b, a) = 1 -^* \widehat{\text{Sd}}(a, b)$.
- (2) $\text{Sd}(a, c) = \widehat{\text{Sd}}(a, b) +^* \widehat{\text{Sd}}(b, c)$ modulo $\mathbb{Z} \cup \mathbb{Z}^*$, that is, $\text{Sd}(a, b) +^* \text{Sd}(b, c) -^* \text{Sd}(a, c) \subset \mathbb{Z}^*$.

By (1), Sd is not symmetric, that is, for $a, b \in M$, $\text{Sd}(a, b) \neq \text{Sd}(b, a)$ and so it is called a directed distance.

Now we assign each 1-simplex f a value n_f in $\mathbb{R} \cup \mathbb{Q}^*$ as follows : There are $a, b \in M$ such that $[a, b] = f$, and we define n_f as $\text{Sd}(a, b)$. Then n_f is well-define, that is, it does not depend on the choice of a, b because if $a_i, b_i \in M$ satisfy $[a_0, b_0] = [a_1, b_1] = f$, then $a_0 b_0 \equiv a_1 b_1$ and $\text{Sd}(a_0, b_0) = \text{Sd}(a_1, b_1)$. We also assign each 1-shell $s = f_{01} + f_{12} - f_{02}$ to a multivalue n_s in $\mathbb{R} \cup \mathbb{Q}^*$ as follows : $n_s = n_{f_{01}} +^* n_{f_{12}} -^* n_{f_{02}}$. This value is also related with the distance of end points. Let (a, a') be an endpoint pair of s , then $\text{Sd}(a, a') = n_s$ modulo $\mathbb{Z}^*$. Using this assignment of 1-shells, we give a necessary and sufficient condition for a 1-shell to be a boundary of a 2-chain :

Theorem 3.8. *A 1-shell $s = f_{12} - f_{02} + f_{01}$ is a boundary of a 2-chain in p if and only if*

$$n_s = n_{01} +^* n_{12} +^* n_{20} \subset \mathbb{Z}^*,$$

where $n_{01} = n_{f_{01}}$, $n_{12} = n_{f_{12}}$, $n_{20} = -^* n_{f_{02}}$. Moreover it is equivalent to that the two end points of s are Lascar equivalent over $\emptyset$.

Proof. ($\Rightarrow$) Let α be a 2-chain having the boundary s . By Fact 1.14, we may assume $\alpha = \sum_{i=0}^{2n} (-1)^i a_i$ be a chain-walk from f_{01} to $-f_{02}$ with $\text{supp}(\alpha) = \{0, 1, 2\}$. Let $[3] = \{0, 1, 2\}$. From Theorem 2.5, there are independent elements $d_0, d_1, \dots, d_{2n+2}$ such that

- $a_i([3]) \equiv [d_0, d_{2i+1}, d_{2i+2}]_{[3]}$ if i is even, and $a_i([3]) = [d_0, d_{2i+2}, d_{2i+1}]_{[3]}$ if i is odd;
- For some even number $0 \leq i_0 \leq 2n$, $[d_{2i_0+1}, d_{2i_0+2}]_{\{1,2\}} = f_{12}(\{1, 2, \})$; and

- For each even number $0 \leq j_0 \neq i_0 \leq 2n$, there is an odd number $0 \leq j_1 \leq 2n$ such that $[d_{2j_0+1}, d_{2j_0+2}]_{\{1,2\}} = [d_{2j_1+2}, d_{2j_1+1}]_{\{1,2\}}$.

Then $\text{Sd}(d_1, d_0) +^* \text{Sd}(d_0, d_{2n+2}) = -^* n_{01} -^* n_{20}$ and by Fact 3.7 (1), $\text{Sd}(d_1, d_2) +^* \text{Sd}(d_2, d_3) +^* \dots +^* \text{Sd}(d_{2n+1}, d_{2n+2}) = n +^* n_{12}$. By Fact 3.7 (2), $\text{Sd}(d_1, d_{2n+2}) \in (-^* n_{01} -^* n_{20}) \cap (n +^* n_{12})$. Since $\{0\}^* = (-^* n_{01} -^* n_{20}) +^* (n_{01} +^* n_{20})$ and $(n +^* n_{12}) +^* (n_{01} +^* n_{20}) = n +^* (n_{01} +^* n_{12} +^* n_{20})$, these two equations imply $n +^* (n_{01} +^* n_{12} +^* n_{20}) \subset \{0\}^*$. Therefore, $n_{01} +^* n_{12} +^* n_{20} \subset \{0\}^* -^* \{n\}^* = \{-n\}^*$ for $n \in \mathbb{N} \subset \mathbb{Z}$.

($\Leftarrow$) Suppose $n_{01} +^* n_{12} +^* n_{20} \subset \{n\}^*$ for some $n \in \mathbb{Z}$. Then there are independent elements a, b, c, a' such that

$$[a, b]_{\{0,1\}} = f_{01}(\{0, 1\}), [b, c]_{\{1,2\}} = f_{12}(\{1, 2\}), [a', c]_{\{0,2\}} = f_{02}(\{0, 2\}).$$

So, $\text{Sd}(a, b) = n_{01}$, $\text{Sd}(b, c) = n_{12}$, $\text{Sd}(c, a) = n_{20}$, and $\text{Sd}(a, a') \in n_{01} +^* n_{12} +^* n_{20}$. Thus $\text{Sd}(a, a') \in \{n\}^*$ and $\text{Sd}(a, a') \in \{0\}^*$. Since $\{a, a'\}$ is independent, $\text{Sd}(a, a') \in \{0\}^* \setminus \{0\}$, that is, $\text{Sd}(a, a') = \epsilon$ or $\text{Sd}(a, a') = 1 - \epsilon$.

We will find $d \in M$ such that $a \equiv_d a'$ and $d \perp_{abca'}$. Consider a partial type $\Sigma(x) = \{s < \text{Sd}(x, a) < t \leftrightarrow s < \text{Sd}(x, a') < t\}_{s < t \in [0,1]}$. Consider finitely many pairs (s_i, t_i) with $s_i < t_i$ and a formula

$$\bigwedge (s_i < \text{Sd}(x, a) < t_i \leftrightarrow s_i < \text{Sd}(x, a') < t_i).$$

We may assume $s_i \leq s_0 < t_0 \leq t_i$. It is enough to show that

$$s_0 < \text{Sd}(x, a) < t_0 \leftrightarrow s_0 < \text{Sd}(x, a') < t_0$$

is satisfiable. Suppose $s_0 < \text{Sd}(x, a) < t_0$ is satisfiable. Then there is a pair (s, t) such that $s_0 < s < t < t_0$ and $s < \text{Sd}(x, a) < t$ is satisfiable. Let $e \in M$ be independent from a such that $s < \text{Sd}(e, a) < t$ holds. Since $\text{Sd}(a, a') \in \{0\}^* \setminus \{0\}$, there is a pair (s', t') such that $s' < t'$, $s_0 < s + s' < t + t' < t_0$, and $s' < \text{Sd}(a, a') < t'$. Then, $s < \text{Sd}(e, a) < t$ and $s' < \text{Sd}(a, a') < t'$ imply $s + s' < \text{Sd}(e, a') < t + t'$. Since $s_0 < s + s' < t + t' < t_0$, $s_0 < \text{Sd}(e, a') < t_0$ and $s_0 < \text{Sd}(x, a') < t_0$ is satisfiable. By the same way, $s_0 < \text{Sd}(x, a') < t_0 \rightarrow s_0 < \text{Sd}(x, a) < t_0$.

Therefore, there is $d \in M$ such that $\Sigma(d)$ and $\text{Sd}(d, a) = \text{Sd}(d, a')$. Moreover we may assume that $\{a, b, c, a', d\}$ is independent by taking $d \perp_{aa'} bc$. Then, there is a 2-chain $\alpha = a_0 + a_1 - a_2$, where

- $\text{supp}(a_0) = \{0, 1, 3\}$, $\text{supp}(a_1) = \{1, 2, 3\}$, and $\text{supp}(a_2) = \{0, 2, 3\}$;
- $a_0(\{0, 1, 3\}) = [a, b, d]_{\{0,1,3\}}$, $a_1(\{1, 2, 3\}) = [b, c, d]_{\{1,2,3\}}$, and $a_2(\{0, 2, 3\}) = [a', c, d]_{\{0,2,3\}}$;
- $a_0^{01} = f_{01}$, $a_1^{12} = f_{12}$, and $a_2^{02} = f_{02}$; and
- $a_0^{03} = a_2^{03}$, $a_0^{13} = a_1^{13}$, and $a_1^{23} = a_2^{23}$.

Then $\partial\alpha = f_{0,1} + f_{1,2} - f_{02} + (a_2^{03} - a_0^{03})$ and $\partial\alpha = f_{0,1} + f_{1,2} - f_{02}$.

Now we show moreover part. Let a, a' be end points of s . If $a \equiv^L a'$, then s is a boundary of 2-chain and $n_s \subset \{n\}^*$ for some $n \in \mathbb{Z}$. Conversely, we assume that $n_s \subset \{n\}^*$ for some $n \in \mathbb{Z}$. In the proof of right-to-left, we found $d \in M$ such that $a \equiv_d a'$. Consider a substructure generated by d . $\text{cl}(d) = \text{dcl}(c) = \text{acl}(c)$. Then $a \equiv_{\text{cl}(d)} a'$ and $a \equiv^L a'$. $\square$

Now we are ready to prove Theorem 3.7. Define a map $\Phi : H_1(p_0) \rightarrow (\mathbb{R} \cup \mathbb{Q}^*)/\mathbb{Z}^*$ as sending $[s]$ into $n_s +^* \mathbb{Z}^*$, and $(\mathbb{R} \cup \mathbb{Q}^*)/\mathbb{Z}^* \cong \mathbb{R}/\mathbb{Z}$ in Appendix B. It is easy to see this map is surjective. Since for an endpoint pair (a, b) of s , $n_s +^* \mathbb{Z}^* = \text{Sd}(a, b) +^* \mathbb{Z}^*$, this map Φ depends on the endpoint pairs of 1-shells. By Theorem 2.2, given 1-shells s_0 and s_1 and endpoint pairs (a, b) and (b, c) with respect to s_0 and s_1 , there is an 1-shell s such that $[s] = [s_0] + [s_1]$ and (a, c) is an endpoint pair of s , thus this map Φ is a group homomorphism. Moreover by Theorem 3.8, it is injective, and therefore it is an isomorphism.

Therefore, we show that the first homology group in p_0 is isomorphic to $\mathbb{R}/\mathbb{Z}$, and it is interesting that from Theorem 3.8, this first homology group is exactly isomorphic to the Lascar group $\text{Gal}_L(\mathcal{M}_1; \emptyset)$.

3.3. Computation of H_1 in $\mathcal{M}_2$. In this subsection, we compute the first homology group of q_0 . Since q_0 is over $\text{acl}^{eq}(\emptyset)$, we work in $\mathcal{M}_2^{eq}$ with constant elements in $\text{acl}^{eq}(\emptyset)$. Since each elements in M^n/E_n is already in $\text{acl}^{eq}(\emptyset)$, and by Theorem 3.4, we may work in $(\mathcal{M}'_2)^{eq} = \mathcal{N}_2^{eq}$. But already noted in the proof of 3.4 (2), the ternary relation $S(x, y, z)$ is definable in $\mathcal{N}_2$ and thus we work in $\mathcal{M}_1^{eq}$. So, by the previous subsection, the first homology group of q_0 is same with one of p_0 and it is isomorphic to $\mathbb{R}/\mathbb{Z}$, which is also Lascar group over $\text{acl}^{eq}(\emptyset)$ in $\mathcal{M}_2$.

We conjecture that there are only automorphisms described in Theorem 2.4 in the kernel of the canonical epimorphism in Theorem 2.3.

Question 3.8. Let $T = T^{eq}$ be a rosy theory, and $\mathcal{C} \models T$. For $A = \text{acl}(A)$, for a strong type p over A , let $\Psi : \text{Aut}_A(\mathcal{C}) \rightarrow H_1(p)$ be a canonical epimorphism. Then, is the kernel of Ψ exactly generated by automorphisms in the following :

- (1) $\text{Aut}_{\text{acl}(aA)}(\mathcal{C})$ for $a \models p$;
- (2) $\text{Autf}(\mathcal{C})$; and
- (3) $(\Psi')^{-1}([G, G])$, where G and Ψ' are in Theorem 2.4,

so, is $H_1(p) \cong (\text{Gal}_L(\mathcal{C}; A) / \langle \bar{\sigma} : \sigma \in \text{Aut}_{\text{acl}(Aa)}, a \models p \rangle)^{ab}$? Furthermore if any algebraic closure is again a substructure, then is $H_1(p) \cong \text{Gal}_L(\mathcal{C}; A)^{ab}$?

Fortunately, the answer for Question 3.8 is yes for known examples in [4][8] and our two examples.

4. APPENDIX

4.1. Appendix A. We show the possible number of bounded type-definable equivalence classes on a strong type is 1 or at least $2^{\aleph_0}$. Let $T(= T^{eq})$ be any theory of a language $\mathcal{L}$ and let $\mathcal{C}$ be a monster model of T . Fix a small subset, $A = \text{acl}(A)$ and choose a strong type $p(\bar{x})$ over A (with $\bar{x}$ of possibly infinite length). We shall denote $\bar{x}$ as x conventionally.

Theorem 4.1. *Let $E(x, y)$ be a bounded A -type-definable equivalence relation on $p(x)$ and denote p/E for the set of E -classes on p . Then,*

$$|p/E| = 1 \text{ or } \geq 2^{\aleph_0}.$$

Proof. For a convention, we assume $A = \emptyset$. We divide two cases that p/E is finite and p/E is infinite.

Case 1. p/E is finite : Suppose p/E is finite. Let $a_0, \dots, a_n \models p$ be representatives of all distinct classes in p/E , and let $\bar{a} = (a_0, a_1, \dots, a_n)$. At first, we show that E is relatively definable on p . Consider two type-definable formula $E(x, a_0)$ and $\bigvee_{i>0} E(x, a_i)$ partitioning p , and by compactness, $p(x) \models E(x, a_0) \leftrightarrow \phi(x, a_0)$ for some formula $\phi(x, z)$ such that $E(x, a_0) \models \phi(x; a_0)$. Since $a_0 \equiv a_i$, $p(x) \models E(x, a_i) \leftrightarrow \phi(x, a_i)$ for all $i \leq n$. Thus, $p(x) \wedge p(y) \models E(x, y) \leftrightarrow \psi(x, y; \bar{a})$, where $\psi(x, y; \bar{z}) = \bigvee_i [\phi(x, z_i) \wedge \bigvee_{j \neq i} \phi(y, z_j)]$. Since E is invariant, $p(x) \wedge p(y) \wedge \psi(x, y; \bar{z}) \wedge \text{tp}(\bar{a})(\bar{z}) \models \psi(x, y; \bar{a}) (\leftrightarrow E(x, y))$. By compactness, there is a formula $\psi'(\bar{z})$ in $\text{tp}(\bar{a})(\bar{z})$ such that $p(x) \wedge p(y) \wedge \psi(x, y; \bar{z}) \wedge \psi'(\bar{z}) \models \psi(x, y; \bar{a})$. Take $\theta(x, y) \equiv \exists \bar{z} (\psi'(\bar{z}) \wedge \psi(x, y; \bar{z}))$. Then $p(x) \wedge p(y) \models \theta(x, y) \leftrightarrow \psi(x, y; \bar{a})$. Therefore E is relatively definable on p by the formula θ . Moreover, we may assume $\theta(x, y)$ is a reflexive and symmetric relation by taking $x = y \vee (\theta(x, y) \wedge \theta(y, x))$.

Next, we find a finite $\emptyset$ -definable equivalence relation E' such that $p(x) \wedge p(y) \models E(x, y) \leftrightarrow E'(x, y)$. Since E is an equivalence relation,

$$\begin{aligned} p(x) \wedge p(y) \wedge p(z) &\models \bigvee_i \theta(x, a_i) \wedge \bigvee_i \theta(y, a_i) \wedge \bigvee_i \theta(z, a_i) \\ &\wedge \bigwedge_i (\theta(x, a_i) \rightarrow \bigwedge_{i \neq j} \neg \theta(x, a_j)) \\ &\wedge \bigwedge_i (\theta(y, a_i) \rightarrow \bigwedge_{i \neq j} \neg \theta(y, a_j)) \\ &\wedge \bigwedge_i (\theta(z, a_i) \rightarrow \bigwedge_{i \neq j} \neg \theta(z, a_j)) \\ &\wedge (\theta(x, y) \wedge \theta(y, z) \rightarrow \theta(x, z)). \quad (*) \end{aligned}$$

Again by compactness, there is $\delta(x) \in p(x)$ such that

$$\delta(x) \wedge \delta(y) \wedge \delta(z) \models (*).$$

Define a definable equivalence relation $E'(x, y) \equiv [\neg \delta(x) \wedge \neg \delta(y)] \vee [\delta(x) \wedge \delta(y) \wedge \forall z (\delta(z) \rightarrow (\theta(z, x) \leftrightarrow \theta(z, y)))]$.

Claim 4.1. *The equivalence relation E' is finite.*

Proof. First, $\neg \delta(x)$ is a E' -class. We show that on δ , the E' -classes are of the form of $\theta(x, a_i) \wedge \delta(x)$. By the choice of δ , it is partitioned by $\{\theta(x, a_i) \wedge \delta(x)\}_{i \leq n}$.

1) We show that $\models \theta(x, a_i) \wedge \delta(x) \rightarrow E'(x, a_i)$: Choose $b \models \theta(x, a_i) \wedge \delta(x)$. Take $c \models \delta(x) \wedge \theta(x, a_i)$. Since θ is transitive on δ and $\theta(b, a_i)$ holds, $\theta(c, b)$ holds. Conversely, if $d \models \delta(x) \wedge \theta(x, b)$, then by transitivity of θ on δ , $\theta(d, a_i)$ holds. Therefore, $E'(b, a_i)$ holds.

2) For $i \neq j$, $\neg E'(a_i, a_j)$: Suppose for some $i \neq j$, $E'(a_i, a_j)$ holds. Then $\theta(a_i, a_j)$ holds but it is impossible since $a_i, a_j \models p$ and θ and E are same on $p \times p$.

By 1) and 2), the E' -classes are of the form of $\theta(x, a_i) \wedge \delta(x)$ or $\neg \delta(x)$ and E' is a finite equivalence relation. $\square$

From the proof of Claim 4.1, E' and E are the same equivalence relation on $p \times p$. Since E' is finite and p is a strong type, $p/E = p/E'$ and there are only one E -class in p .

Case 2. p/E is infinite : Suppose p/E is infinite. Let $\kappa = |p/E|$. If E is definable, then by compactness, $|p/E| \geq \kappa'$ for any $\kappa' < |\mathcal{C}|$ and E is not bounded. So E is type-definable and $E(x, y) \equiv \bigwedge_{i < \lambda} \phi_i(x, y)$, where $\phi_i(x, y)$ is a formula and λ is an infinite cardinal. We may assume $\phi_i(x, y)$ is reflexive and symmetric by taking $x = y \vee (\phi_i(x, y) \wedge \phi_i(y, x))$

instead of $\phi_i(x, y)$ for each $i < \lambda$, and that for $i < j < \lambda$, $\models \phi_j(x, y) \rightarrow \phi_i(x, y)$ ($\dagger$) by taking $\phi_j(x, y) \wedge \phi_i(x, y)$. Moreover, by compactness, we may assume that for $\models \exists z(\phi_{i+1}(x, z) \wedge \phi_{i+1}(z, y)) \rightarrow \phi_i(x, y)$ ($\ddagger$). Let $\{a_k \models p\}_{k < \kappa}$ be the set of representatives of E -classes.

Claim 4.2. *For each $i < \lambda$ and $k < \kappa$, $\phi_i(x, a_k)(\mathcal{C})$ contains infinitely many E -classes.*

Proof. Fix $i < \lambda$. By compactness, there are finitely many $k_0 < k_1 < \dots < k_n$ such that $p \models \bigvee_j \phi_i(x, a_{k_j})$. By Pigeonhole Principle, some $\phi_i(x, a_{k_l})$ contains infinitely many a_k 's. By ($\dagger$) and ($\ddagger$), $\phi_i(x, a_{k_l})$ contains infinitely many E -classes. Since $a_n \equiv a_m$ for $n, m < \kappa$ and E is invariant, each $\phi_i(x, a_k)$ contains infinitely many E -classes. $\square$

Claim 4.3. *For each $i < \lambda$ and $k < \kappa$, there are $i < j < \lambda$ and $k_0, k_1 < \kappa$ such that*

$$\models [(\phi_j(x, a_{k_0}) \vee \phi_j(x, a_{k_1})) \rightarrow \phi_i(x, a_k)] \wedge [\neg \exists x(\phi_j(x, a_{k_0}) \wedge \phi_j(x, a_{k_1}))].$$

Proof. Fix $i < \lambda$ and $k < \kappa$. By Claim 4.2, $\phi_i(x, a_k)$ contains infinitely many E -classes. Choose two E -classes in $\phi_i(x, a_k)$ and let a_{k_0} and a_{k_1} be representatives of two classes respectively. Since $E(x, a_{k_0})(\mathcal{C})$ and $E(x, a_{k_1})(\mathcal{C})$ are disjoint, by compactness, for some $j > i$, $\phi_j(x, a_{k_0})(\mathcal{C})$ and $\phi_j(x, a_{k_1})(\mathcal{C})$ are disjoint and we are done. $\square$

From Claim 4.2, 4.3 and the fact that the cofinality of λ is at least $\aleph_0$, we get a binary tree $\mathcal{B} : 2^{<\omega} \rightarrow \omega \times \kappa$ such that for each $b \in 2^{<\omega}$, $\mathcal{B}(b0) = (j, k_0)$ and $\mathcal{B}(b1) = (j, k_1)$ where if $\mathcal{B}(b) = (i, k)$, then $j < \omega$ and $k_0, k_1 < \kappa$ satisfies Claim 4.3 for (i, k) . Then for each $\tau \in 2^\omega$, we get a set of formula $\{\phi_{i(\tau \upharpoonright n)}(x, a_{k(\tau \upharpoonright n)})\}$, where $\mathcal{B}(\tau \upharpoonright n) = (i(\tau \upharpoonright n), k(\tau \upharpoonright n))$ for each $n \in \omega$. By the choice of $\mathcal{B}$, for $\tau_0 \neq \tau_1 \in 2^\omega$, $\bigcap_n \phi_{i(\tau_0 \upharpoonright n)}(x, a_{k(\tau_0 \upharpoonright n)})(\mathcal{C})$ and $\bigcap_n \phi_{i(\tau_1 \upharpoonright n)}(x, a_{k(\tau_1 \upharpoonright n)})(\mathcal{C})$ are disjoint, and each contains E -classes. Thus, p/E has at least $2^{\aleph_0}$ many elements. $\square$

4.2. Appendix B. We see how to recover a real ordered group $(\mathbb{R}, +)$ from a dense linear order extending $(\mathbb{Q}, <)$ using Dedekind cut. Consider a language $\mathcal{L}_{od, \mathbb{Q}} = \{<\} \cup \{r\}_{r \in \mathbb{Q}}$ and a $\mathcal{L}_{od, \mathbb{Q}}$ -structure $\mathcal{U} = (U, <, r : r \in \mathbb{Q})$ which is a saturated dense linear order extending $(\mathbb{Q}, <)$. Then $\text{Th}(\mathcal{U})$ has quantifier elimination.

Consider the 1-types over empty set, $S_1(\emptyset)(= S_1)$. Then by quantifier elimination, any 1-type p has one of the following forms : For $r \in \mathbb{Q}$ and $r' \in \mathbb{R} \setminus \mathbb{Q}$,

- (1) $\{x = r\}$;
- (2) $\{l < x < r \mid l < r\}$;
- (3) $\{r < x < u \mid r < u\}$; and

$$(4) \{l < x < u \mid l < r' < u\}.$$

For a subset $S \subset \mathbb{Q}$, we write $S^* := S \cup \{s \pm \epsilon \mid s \in S\}$, where we consider ϵ as infinitesimal. So we can identify S_1 with the set $\mathbb{R} \cup \mathbb{Q}^*$ in the following way : For $r \in \mathbb{Q}$ and $r' \in \mathbb{R} \setminus \mathbb{Q}$,

- (1) $\{x = r\} \leftrightarrow r$;
- (2) $\{l < x < r \mid l < r\} \leftrightarrow (r - \epsilon)$;
- (3) $\{r < x < u \mid r < u\} \leftrightarrow (r + \epsilon)$; and
- (4) $\{l < x < u \mid l < r' < u\} \leftrightarrow r'$.

Next we define a group-like structure on S_1 . Define a plus-like operation $+^* : S_1 \times S_1 \rightarrow \mathcal{P}(S_1)$ as follows :

$$p_1 +^* p_2 := \{p \mid p \models (l_1 + l_2 < x < u_1 + u_2), p_i \models l_i < x < u_i\},$$

and define a minus-like operation $-^* : S_1 \rightarrow S_1$ as follows :

$$(-^* p) := \{-u < x < -l \mid p \models l < x < u\}.$$

We define a composition of plus operation as follows :

$$(p_1 +^* p_2) +^* p_3 := \bigcup_{p \in p_1 +^* p_2} p +^* p_3,$$

and

$$p_1 +^* (p_2 +^* p_3) := \bigcup_{p \in p_2 +^* p_3} p_1 +^* p.$$

Then $+^*$ and $-^*$ is commutative, associate, and distributive. And for any $p_1, \dots, p_k \in S_1$ and $k \geq 1$,

$$|p_1 +^* \dots +^* p_k| \leq 3$$

We write $p_1 -^* p_2$ for $p_1 +^* (-^* p_2)$. These two notions are naturally assigned to $\mathbb{R} \cup \mathbb{Q}^*$ and they are defined as follows :

- (1) (a) If both r_1 and r_2 are in $\mathbb{R}$ and let $r = r_1 + r_2$, then

$$r_1 +^* r_2 := \begin{cases} r & \text{if } \{r\} \in \mathbb{R} \setminus \mathbb{Q} \\ r & \text{if } \{r\} \in \mathbb{Q} \text{ and } r_1, r_2 \in \mathbb{Q} \\ \{r - \epsilon, r, r + \epsilon\} & \text{if } r \in \mathbb{Q} \text{ and } r_1, r_2 \notin \mathbb{Q} \end{cases}$$

;

- (b) If $r_1 \in \mathbb{R} \setminus \mathbb{Q}$ and $r_2 = q \pm \epsilon \in \mathbb{Q}^*$, then $r_1 +^* r_2 := \{r_1 + q\}$;
- (c) If $r_1 \in \mathbb{Q}$ and $r_2 = q \pm \epsilon \in \mathbb{Q}^*$, then $r_1 +^* r_2 := \{(r_1 + q) \pm \epsilon\}$;
- (d) If $r_1 = p \pm \epsilon$ and $r_2 = q \pm \epsilon \in \mathbb{Q}^*$, then $r_1 +^* r_2 := \{(p + q) \pm \epsilon\}$;
- (e) If $r_1 = p \pm \epsilon$ and $r_2 = q \mp \epsilon \in \mathbb{Q}^*$, then $r_1 +^* r_2 := \{(p + q) - \epsilon, (p + q), (p + q) + \epsilon\}$.
- (2) (a) If $r_1 \in \mathbb{R}$, then $-^* r_1 := -r_1$;
- (b) If $r_1 = p \pm \epsilon \in \mathbb{Q}^*$, then $-^* r_1 := -p \mp \epsilon$.

Now we induce a group structure from $(S_1, +^*, -^*)$. Define an equivalence relation $\equiv_0$ on S_1 ,

$$p_1 \equiv_0 p_2 \text{ iff } p_1 -^* p_2 \subset \{0 - \epsilon, 0, 0 + \epsilon\},$$

and denote $[p]_0$ for $p \in S_1$ for the equivalence class. Since $\{0 - \epsilon, 0, 0 + \epsilon\}$ is closed under $+^*$ and $-^*$, $+^*$ and $-^*$ are extended on $S_1 / \equiv_0$. Then $(S_1 / \equiv_0, +^*, -^*, [\text{tp}(0)]_0)$ is a group. Actually it is isomorphic to $(\mathbb{R}, +, -, 0)$.

Theorem 4.2. $(S_1 / \equiv_0, +^*, -^*, [\text{tp}(0)]_0) \cong (\mathbb{R}, +, -, 0)$.

And define an equivalence relation $\equiv_{\mathbb{Z}}$ on S_1 ,

$$p_1 \equiv_{\mathbb{Z}} p_2 \text{ iff } p_1 -^* p_2 \subset \mathbb{Z}^*,$$

and denote $[p]_{\mathbb{Z}}$ for the equivalence class. As same as $\equiv_0$, $\mathbb{Z}^*$ is closed under $+^*$ and $-^*$ and $+^*$ and $-^*$ are extended on $S_1 / \equiv_{\mathbb{Z}}$. And $(S_1 / \equiv_{\mathbb{Z}}, +^*, -^*, [\text{tp}(0)]_{\mathbb{Z}})$ is isomorphic to $(\mathbb{R}/\mathbb{Z}, +, -, 0)$ as groups.

Theorem 4.3. $(S_1 / \equiv_{\mathbb{Z}}, +^*, -^*, [0]_{\mathbb{Z}}) \cong (\mathbb{R}/\mathbb{Z}, +, -, 0)$.

These two equivalences $\equiv_0$ and $\equiv_{\mathbb{Z}}$ are defined on $\mathbb{R} \cup \mathbb{Q}^*$ and

$$(\mathbb{R} \cup \mathbb{Q}^*) / \equiv_0 \cong \mathbb{R} \text{ and } (\mathbb{R} \cup \mathbb{Q}^*) / \equiv_{\mathbb{Z}} \cong \mathbb{R}/\mathbb{Z}.$$

Moreover we can define a multiplication-like operation $\times^*$ on S_1 as similar as the plus-like operation $+^*$. Note that for $r_0 \in \mathbb{Q}$ and $r \in \mathbb{R} \cup \mathbb{Q}^*$,

$$\begin{aligned} \text{tp}(r_0) \times^* \text{tp}(r) &:= \begin{cases} \text{tp}(r_0 r) & \text{if } r \in \mathbb{R} \\ \text{tp}(r_0 q + \epsilon) & \text{if } r = q + \epsilon, q \in \mathbb{Q} \\ \text{tp}(r_0 q - \epsilon) & \text{if } r = q - \epsilon, q \in \mathbb{Q} \end{cases} \end{aligned}$$

These plus-, minus-, multiplication-like operations make

$$(S_1 / \equiv_0, [\text{tp}(0)]_0, [\text{tp}(1)]_0, +^*, -^*, \times^*) \cong (\mathbb{R}, 0, 1, +, -, \times),$$

and

$$(S_1 / \equiv_{\mathbb{Z}}, [\text{tp}(0)]_{\mathbb{Z}}, [\text{tp}(1)]_{\mathbb{Z}}, +^*, -^*, \times^*) \cong (\mathbb{R}/\mathbb{Z}, 0, 1, +, -, \times).$$

REFERENCES

- [1] Enrique Casanovas, Daniel Lascar, Anand Pillay, and Martin Ziegler. Galois groups of first order theories. *Journal of Mathematical Logic*, **1** (2001) 305-319.
- [2] Clifton Ealy and Alf Onshuus. Characterizing rosy theories. *Journal of Symbolic Logic*, **72** (2007) 919-940.
- [3] John Goodrick, Byunghan Kim, and Alexei Kolesnikov. Homology groups of types in model theory and the computation of $H_2(p)$. *Journal of Symbolic Logic*, **78** (2013), pp. 1086-1114.
- [4] John Goodrick, Byunghan Kim, and Alexei Kolesnikov. Amalgamation functors and homology groups in model theory. will appear in the proceedings of ICM 2014.

- [5] John Goodrick, Byunghan Kim, and Alexei Kolesnikov. Homology groups of types in stable theories and the Hurewicz correspondence. Submitted.
- [6] Byunghan Kim. *Simplicity Theory*, Oxford University Press, (2014).
- [7] Byunghan Kim, Alexei Kolesnikov, and Akito Tsuboi. Generalized amalgamation and n -simplicity. *Annals of Pure and Applied Logic*, **155** (2008), 97-114
- [8] Byunghan Kim, SunYoung Kim, and Junguk Lee, A classification of 2-chains having 1-shell boundaries in rosy theories, *Journal of Symbolic Logic*, **80**, (2015), 322-340.
- [9] SunYoung Kim, and Junguk Lee. More on 2-chains with 1-shell boundaries in rosy theories. accepted in *Journal of the mathematical society of Japan*.
- [10] Krzysztof Krupinski, and Tomasz Rzepecki. Smoothness of bounded invariant equivalence relations. accepted in *Journal of Symbolic Logic*.
- [11] Ludomir Newelski. The diameter of a Lascar strong type. *Fundamenta Mathematicae*, **176** (2003), 157-170.
- [12] Bruno Poizat. *A course in model theory: an introduction to contemporary mathematical logic*, Springer, (2000).

DEPARTMENT OF MATHEMATICS, YONSEI UNIVERSITY, 50 YONSEI-RO, SEODAEMUN-GU, SEOUL 120-749, KOREA

E-mail address: ljw@yonsei.ac.kr